



ETILHEXILGLICERINA

INCI Name: Ethylhexylglycerin

A etilhexilglicerina é um derivado sintético do glicerol, amplamente utilizado em cosméticos como conservante auxiliar e agente antimicrobiano suave. **Uso Externo. Importante: Produto exclusivamente para aplicações cosméticas.** Mantenha o produto fora do alcance de crianças e ao abrigo da luz e calor. Evite contato com os olhos e boca. Em caso de irritação, retire imediatamente o produto com água corrente, suspenda o uso e procure orientação de um profissional da área da saúde.



Distribuído por Engenharia das Essências
Comercial Ltda. Rua Ronaldo de Carvalho, 292
Jd. São Paulo/SP - CEP 02043-000.
CNPJ 36.437.409/0001-17 - Ind.Brasileira.
Envasado por CNPJ: 04.556.017/0001-80
AFE: 2.05.921-1. SAC 11 2978-6311.

www.engenhariadasessencias.com

☐ ☐
500g 1000g

the 1990s, the incidence of *S. flexneri* infections in the United Kingdom has increased, and the incidence of *S. flexneri* infection in the United States has increased in the 1980s and 1990s [10].

There is a paucity of data on the incidence of *S. flexneri* infection in the United Kingdom. In the 1980s, *S. flexneri* was the most commonly isolated serotype of *Shigella* from patients with shigellosis in the United Kingdom [11]. In the 1990s, *S. flexneri* was the most commonly isolated serotype of *Shigella* from patients with shigellosis in the United Kingdom [12].

The purpose of this study was to determine the incidence of *S. flexneri* infection in the United Kingdom in the 1990s. The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection is high, and the incidence of *S. flexneri* infection is high in the United States.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection is high, and the incidence of *S. flexneri* infection is high in the United States.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection is high, and the incidence of *S. flexneri* infection is high in the United States.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection is high, and the incidence of *S. flexneri* infection is high in the United States.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection is high, and the incidence of *S. flexneri* infection is high in the United States.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection is high, and the incidence of *S. flexneri* infection is high in the United States.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection is high, and the incidence of *S. flexneri* infection is high in the United States.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection is high, and the incidence of *S. flexneri* infection is high in the United States.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection is high, and the incidence of *S. flexneri* infection is high in the United States.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection is high, and the incidence of *S. flexneri* infection is high in the United States.

The study was conducted in the United Kingdom, where the incidence of *S. flexneri* infection is high, and the incidence of *S. flexneri* infection is high in the United States.